



ISSN 2348 - 0319

Journal home page: <http://www.journalijar.com>

INTERNATIONAL JOURNAL  
OF INNOVATIVE AND  
APPLIED RESEARCH

## RESEARCH ARTICLE

## FRACTIONAL FREE ELECTRON LASER EQUATION AND M-SERIES

Manoj Sharma

1 Department of Mathematics RJIT, BSF Academy, Tekanpur.

**Abstract:**

In recent year's fractional free electron laser (FEL) equation are studied due to their usefulness and importance in mathematical physics, The aim of present paper is to find the solution of generalized fractional order free electron laser (FEL) equation, using M-SERIES. The results obtained here is moderately universal in nature. Special cases, relating to the exponential function is also considered.

**Key Words:** Fractional free electron laser (FEL) equation, M-SERIES, Riemann-Liouville operator.

**1. Introduction:****The Fractional Free Electron Laser Equation:**

The unsaturated behaviour of the free electron laser (FEL) is governed by the following first order integro differential equation of Volterra – type [3,4]. :

$$D_T a(T) = -i\pi g_0 \int_0^T \xi a(T - \xi) e^{iv\xi} d\xi, \quad 0 \leq T < 1 \quad \dots(1.1)$$

where T is a dimensionless time variable,  $g_0$  is a positive constant known as the small–signal gain and the constant  $v$  is the detuning parameter. The functional (T) is a complex-field amplitude which is assumed to be dimensionless and satisfies the initial condition  $a(0) = 1$ . Here we employ the Riemann-Liouville definition of fractional integral equation defined by a simplified version of (1.1) changing the scale by putting  $t = x\sigma$  and  $a = 0$  this yields

$$R_x^\alpha f(x) \equiv I_x^\alpha f(x) = \frac{x^\alpha}{\Gamma(\alpha)} \int_0^1 (1-\sigma)^{\alpha-1} f(x\sigma) d\sigma, \quad \text{Re } \sigma \geq 0 \quad \dots(1.2)$$

The definition (1.2) can be written as

$$R_x^\alpha f(x) \equiv I_x^\alpha f(x) = \frac{d^n}{dx^n} R_x^{\alpha+n} f(x), \quad \text{Re } (\alpha + n) > 0 \quad \dots(1.3)$$

Boyardjiev et al. [3] have treated a non homogeneous case of (1.2) in which the ordinary first derivative  $D_T$  is replaced by the fractional  $D_T^\alpha$  with  $\alpha > 0$ , that is

$$D_T^\alpha a(T) = \lambda \int_0^T t a(T-t) e^{ivt} dt + \beta e^{ivt}, \quad 0 \leq T \leq 1 \quad \dots(1.4)$$

with  $\beta, \lambda, \in \mathbb{C}$  and  $v \in \mathbb{R}$ . Furthermore the following generalization of (1.4) has been considered by the authors [2]

$$D_T^\alpha a(T) = \lambda \int_0^T t^\delta a(T-t) e^{ivt} dt + \beta e^{ivt}, \quad 0 \leq T \leq 1 \quad \dots(1.5)$$

where  $\beta, \lambda, \in \mathbb{C}, v \in \mathbb{R}$  and  $\delta > -1$ , In the present section, we investigate a further generalization of equation (1.5), whereby the exponential term is replaced by the M-SERIES

**2. The Generalized Equation:**

The generalization of equation (1.5) obtained by replacing  $e^{ivt}$  by  ${}_p M_q^\alpha(z)$  takes the form

$${}_p M_q^\alpha(z) = \sum_{k=0}^{\infty} \frac{(a_1)_k \dots (a_p)_k}{(b_1)_k \dots (b_q)_k} \frac{z^k}{\Gamma(\alpha k + 1)}$$

$$D_T^\alpha a(T) = \lambda \int_0^T t^\delta a(T-t) {}_p M_q^\alpha(ivt) dt + \beta {}_p M_q^\alpha(ivt), \quad \dots(2.1)$$

$$0 \leq T \leq 1.$$

where  $\alpha, \beta, \lambda \in \mathbb{C}; v \in \mathbb{R}, \alpha > 0, \delta > -1$ , Given real numbers  $b_k$ , the corresponding initial conditions are :

$$D_T^{\alpha-k} a(T) \Big|_{T=0} = b_k, \quad k = 1, 2, 3, \dots, N \quad \dots(2.2)$$

with  $N = [\alpha] + 1$ , so that  $N - 1 \leq \alpha < N$ . Equation (1.5) can be deduced from (2.1) by taking  $\alpha = 1$  while (2.4) follows when  $\alpha = 1, \delta = 1$  To obtain the solution of (2.1) for the given initial conditions (2.2), we use (2.2).

Let  $\xi = T - t$  in (2.1), so that  $D_T^\alpha a(T) = \lambda \int_0^T (T - \xi)^\delta a(\xi) {}_p M_q^\alpha(iv(t - \xi)) d\xi + \beta {}_p M_q^\alpha(ivT)$

$$\dots(2.3)$$

Using the series representation for  ${}_p M_q^\alpha(z)$  we get

$$a(T) = a_0(T) + \lambda I_T^\alpha \left[ \sum_{k=0}^{\infty} \frac{(a_1)_k \dots (a_p)_k}{(b_1)_k \dots (b_q)_k} \frac{(iv)^k}{\Gamma(\alpha k + 1)} \int_0^T (T - \xi)^{\delta+k} a(\xi) d\xi \right] + \beta I_T^\alpha \left[ \sum_{k=0}^{\infty} \frac{(a_1)_k \dots (a_p)_k}{(b_1)_k \dots (b_q)_k} \frac{(iv)^k}{\Gamma(\alpha k + 1)} \right] \dots(2.4)$$

where  $a_0(T) = \sum_{k=1}^N \frac{b_k}{\Gamma(\alpha - k + 1)} T^{N-k} \quad \dots(2.5)$

By using the Dirichlet formula

$$\int_0^t \int_0^T u(T, s) ds dT = \int_0^t \int_0^t u(T, s) dT ds \quad \dots(2.6)$$

We obtain the following result

$$a(T) = a_0(T) + \frac{\lambda}{\Gamma(\alpha + 1)} \int_0^T a(\xi) (T - \xi)^{\alpha + \delta} {}_p M_q^\alpha (iv(T - \xi)) d\xi + \frac{\beta \Gamma(k + 1) T^\alpha}{\Gamma(\alpha + k + 1)} {}_p M_q^\alpha (ivT) \quad \dots(2.7)$$

Since (2.7) is a Volterra integral equation with continuous kernel, it admits a unique continuous solution (2.4). Finally, we consider some special cases of the generalized fractional integro-differential equation of volterra – type (2.1)

1. If  $\alpha = 1$  and there is no upper and lower parameter

$$D_T^\alpha a(T) = \lambda \int_0^T t^\delta a(T - t) e^{iv(T - \xi)} d\xi + \frac{\beta \Gamma(k + 1) T^\alpha e^{ivT}}{\Gamma(\alpha + k + 1)} \quad \dots(2.8)$$

Equivalently

$$D_T^\alpha a(T) = \lambda \int_0^T t^\delta a(T - t) {}_0F_0(-; -; iv(T - \xi)) + \beta \frac{\Gamma(k + 1)}{\Gamma(\alpha + k + 1)} T^\alpha {}_0F_0(-; -; ivT) \quad \dots(2.9)$$

### 3. Conclusion

In this present work, we have introduced a fractional generalization of the standard free electron laser (FEL) equation. The results of the Advanced generalized fractional free electron laser (FEL) equation and its special case are same as the results of Al-Shammery, A, Kalla, and Khajah, [2](2003).

### References

- [1] Abramowitz, M., Stegun, I. A.: Handbook of Mathematical Functions. Dover (1965), New York.
- [2] Al-Shammery, A. H., Kalla, S.L. and Khajah, H. G.: A fractional generalization of the free electron laser equation. J. FCAA, Vol. 6, No.4 (2003), 501-507.
- [3] Boyadjiev, L., Kalla, S. L. and Khajah, H. G.: Analytical and numerical treatment of a fractional integro-differential equation of Volterra- type Math. Comput. Modelling 25, No. 12 (1997), 1-9.

[4] Dattoli, G., Gianessi, L., Mezi, L., Torre, A. and Caloi, R.: FEL time-evolution operator. Nucl. Instr. Methods A 304 (1991), 541-544.

[5] Dattoli, G., Lorenzutta, S., Maino, G. and Torre, A.: Analytical treatment of the high-gain free electron laser equation Radiat. Phys. Chem. 48, No1 (1996), 29-40.